



Walløe, L., Hjort, N. L., & Thoresen, M. (2019). Start cooling as soon as possible. *Acta Paediatrica*, 108(4), 771.  
<https://doi.org/10.1111/apa.14710>

Early version, also known as pre-print

Link to published version (if available):  
[10.1111/apa.14710](https://doi.org/10.1111/apa.14710)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the submitted manuscript (SM). The accepted author manuscript version (AAM) will be made available online via Wiley at <https://onlinelibrary.wiley.com/doi/full/10.1111/apa.14710>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**Response to:**

**Late Initiation of Therapeutic Hypothermia: Not the New Standard of Care (Bourque SL, Dietz RM.**

**Start Cooling as soon as possible**

Thank you for your supportive comments (1) on our letter (2), regards statistical interpretation of the Laptook et al paper publishing a trial on delayed therapeutic hypothermia (TH) treatment which you supported (3). You also present arguments in favour of allowing this late start of TH after perinatal asphyxia and we would like to comment on those.

You say (1):

‘Laptook and colleagues importantly demonstrated the feasibility and safety of late TH when early initiation of TH may not be possible. Specifically, there were no increased adverse events in the infants who underwent late TH.’

Should one offer treatment with uncertain efficacy as long as the treatment is feasible and has shown no harm? Even if harm was not shown from 96 hours of TH, starting at ~16 hours after birth in the Laptook paper, there are known disadvantages and uncertainties related to hypothermia treatment. The newborn is separated from its parents, and he/she is less likely to receive any breast milk until after rewarming. As an intensive care patient, most infants undergoing TH will be ventilated and sedated, receive antibiotics and parenteral nutrition. We do not know the long-term effects of the necessary sedative drugs delivered during TH (it is unpleasant to be cold as well as ventilated). It is known that drug levels show great individual variability (4), partly due to unpredictable liver and kidney metabolism during cooling. Animal data from different species suggest that opiates damage the immature brain including the cerebellum and starting cooling with a delay of 12 hours in rats after a severe HI insult increased injury (5). These disadvantages need to be weighed against a treatment with disputable effect.

Your next point (1) addresses two clinical problems:

‘Under-recognition of encephalopathy and delivery at centres without cooling capability coupled with prolonged transport time are potential reasons for inability to cool within 6 hours after delivery.’

The first point highlights the need for training of staff and availability of some basic monitoring equipment for cooling like rectal temperature probes for continuous temperature monitoring. Single channel aEEG equipment will document background voltage depression or seizures and help the diagnosis of encephalopathy, a web-camera or smartphone can send pictures and video of the aEEG trace and a neurological examination to the cooling centre. The cooling centre should be able to provide a second opinion fast and advise on starting passive cooling or not. A cohort study showed it was the time of starting cooling and not the time it took to reach 33.5° which is the more important. (6)

We think the effort is better used on early diagnoses with centralised review, cooling during transport and starting cooling within 6 hours after birth.

**References**

1. Bourque SL, Dietz RM. Late initiation of therapeutic hypothermia: not the new standard of care. *Acta Paediatr* 2019; 108: 770
2. Walløe L, Hjort NL, Thoresen M. Major concerns about late hypothermia study. *Acta Paediatrica* 2019; 108: 588-9; e-pub 11 Nov 2018 DOI: 10.1111/apa.14640

3. Bourque SL, Dietz RM. Does late therapeutic hypothermia reduce risk of death or disability? *Acta Paediatr* 2018; 107:1103
4. Frymoyer A, Bonifacio SL, Drover DR, Su F, Wustoff CJ, Van Meurs KP. Decreased morphine clearance in neonates with hypoxic ischemic encephalopathy receiving hypothermia. *J Clin Pharmacol* 2017; 57: 64-76
5. Sabir H, Scull-Brown E, Liu X, Thoresen M - Immediate hypothermia is not neuroprotective after severe hypoxia-ischemia and is deleterious when delayed by 12 hours in neonatal rats. *Stroke* 2012;43:3364-3370 doi: 10.1161/STROKEAHA.112.674481
6. Thoresen M, Tooley J, Liu X, Jary S, Fleming P, Luyt K, Jain A, Cairns P, Harding D, Sabir H - Time Is Brain: Starting Therapeutic Hypothermia within Three Hours after Birth Improves Motor Outcome in Asphyxiated Newborns. *Neonatology* 2013;104:228-233 doi: 10.1159/000353948

Lars Walløe<sup>1</sup>, Nils Lid Hjort<sup>2</sup>, Marianne Thoresen<sup>3</sup>

1. Division of Physiology, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway

2. Division of Statistics and Biostatistics, Department of Mathematics, University of Oslo, Oslo, Norway

3. Neonatal Neuroscience, Translational Health Sciences, BRMS, University of Bristol, Bristol, UK

### **Correspondence**

M Thoresen, Neonatal Neuroscience, Translational Health Sciences, St. Michael's Hospital, University of Bristol, Level D, Southwell Street, Bristol BS2 8EG, UK. Tel: +44 117 342 5226/5607  
Email: [marianne.thoresen@bristol.ac.uk](mailto:marianne.thoresen@bristol.ac.uk)

*Response to:* Late Initiation of Therapeutic Hypothermia: Not the New Standard of Care